

Diabetes Management HEDIS Measures and Tips



Improving Quality Outcomes

Line of Business: Medicaid, Marketplace, Medicare $\bullet \bullet \bullet$

ambetter

BLOOD PRESSURE CONTROL FOR PATIENTS WITH DIABETES (BPD)

Measure Description

The percentage of members 18–75 years of age with diabetes (type 1 and type 2) whose blood pressure was adequately controlled (< 140/90 mm Hg) during the measurement year.

Description	CPT II Codes	
Systolic Blood Pressure less than 130 mm Hg	3074F	
Systolic Blood Pressure 130-139 mm Hg	3075F	
Systolic Blood Pressure greater than or equal to 140 mm Hg	3077F	
Diastolic Blood Pressure less than 80 mm Hg	3078F	
Diastolic Blood Pressure 80-89 mm Hg	3078F	
Diastolic Blood Pressure greater than or equal to 90 mm Hg	3080F	
Remote Patient Monitoring		
Description	CPT Codes	
Remote Blood Pressure Monitoring	93784, 93788, 93790, 99091, 99453, 99454, 99457, 99473, 99474	

Telehealth Visits		
Description	Codes	
Telephonic visit	CPT: 98966-98968, 99441- 99443	
	CPT: 98970-98972, 99421, 99422, 99457	
Online visit	HCPCS: G2012, G0071, G2251, G2252	
Modifier/Place of Services	Modifier : 95 POS: 02, 10	

Codes subject to change

Improving HEDIS Measure

- Patients frequency of visit is based on their blood pressure being controlled. An uncontrolled blood pressure may require more frequency.
- Bill with CPT II codes both systolic and diastolic to report BP frequency for each visit. This can reduce the need for medical record chart reviews.
- Use telehealth when applicable for blood pressure remote patient monitoring and medication management. Patient can only self-report with a digital blood pressure cuff.

Note:

HEDIS captures the last blood pressure reading of the measurement year.



EYE EXAM FOR PATIENTS WITH DIABETES (EED)

Measure Description

The percentage of members 18–75 years of age with diabetes (type 1 and type 2) who had a retinal eye exam.

Description	Codes
Diabetic Eye Exam	CPT: 92002,92004, 92012, 92014,
	92018, 92019, 92134, 92201,
	92202,92227, 92228, 92230, 92235,
	92240, 92250, 92260, 99203- 99205,
	99213- 99215, 99242- 99245
	HCPCS: S0620, S0621 S3000

Description	Codes	
Eye Exam w/ Evidence of Retinopathy	CPT II: 2022F, 2024F, 2026F	
Eye Exam w/o Evidence of Retinopathy	CPT II: 2023F, 2025F, 2033F	
Unilateral Eye Enucleation	CPT: 65091, 65093, 65101, 65103,	
	65105, 65110, 65112, 65114	

Codes subject to change

Improving HEDIS Measure

- Work with a local ophthalmologist or optometrist to establish DRE referral contacts/relationships. Refer to <u>www.pshpgeorgia.com</u> to find-a-provider innetwork
- Educate patients about the difference between a routine eye exam to get new glasses and a comprehensive diabetic eye exam.
- Documentation of hypertensive retinopathy is considered positive for diabetic retinopathy. An annual comprehensive diabetic eye exam is recommended.



Helpful Documentation Tips

- At a minimum, documentation in the medical record must include one of the following:
- A note or letter prepared by an ophthalmologist, optometrist, PCP, or other healthcare professional indicating that an ophthalmoscopic exam was completed by an eye care professional (optometrist or ophthalmologist), the date when the procedure was performed, and the results.
- A chart or photograph indicating the date when the fundus photography was performed and evidence that an optometrist or ophthalmologist reviewed the results. Alternatively, results may be read by a qualified reading center that operates under the direction of a medical director who is a retinal specialist or by a system that provides an artificial intelligence (AI) interpretation.
- Documentation of a negative retinal or dilated eye exam by an optometrist or ophthalmologist in the year prior to the measurement year, results indicating retinopathy was not present.
- Notate anytime in the member's history

GLYCEMIC STATUS ASSESSMENT FOR PATIENTS WITH DIABETES (GSD) Formerly HBD

Measure Description

The percentage of members 18–75 years of age with diabetes (types 1 and 2) whose hemoglobin A1c (HbA1c) was at the following levels during the measurement year:

- Glycemic Status < 8.0%</p>
- Glycemic Status > 9.0%

Description	Codes	
HbA1c Lab Test	CPT: 83036, 83037	
HbA1c – CPT II Codes		
Description	CPT II	
Most recent HbA1c level less than 7.0% (DM)	3044F	
Most recent HbA1c greater than 9.0% (DM)	3046F	
Most recent HbA1c level greater than or equal to 7.0% and less than or equal to 8.0% (DM)	3051F	
Most recent HbA1c level greater than or equal to 8.0% and less than or equal to 9.0% (DM)	3052F	

Codes subject to change

Improving HEDIS Measure

- The frequency of visits should depend on the level of A1c control; members with elevated A1c levels need to be seen more frequently.
- Schedule follow-up visits and A1c testing with diabetic patients to monitor for changes.
- Document the date of the HbA1c with the results.
- Submit the CPT code for the test performed and the CPT II codes to report the A1c results and findings. Also, billing with CPT II codes can reduce medical records chart reviews.

Note

If multiple HbA1c tests were performed in the measurement year, the result from the last test is utilized to close the HEDIS care gap.



Measure Description

The percentage of members 18–85 years of age with diabetes (type 1 and type 2) who received a kidney health evaluation, defined by an estimated glomerular filtration rate (eGFR) and a urine albumin-creatinine ratio (uACR), during the measurement year.

Description	CPT Codes
Estimated Glomerular Filtration Rate (eGFR)	80047, 80048, 80050, 80053,
	80069, 82565
Quantitative Urine Albumin Lab Test	82043
Urine Creatinine Lab Test (uACR)	82570

Codes subject to change

Improving HEDIS Measure

- Routinely refer members with type 1 or type 2 diabetes to a participating lab for an eGFR and uACR.
- Follow up with patients to discuss their lab results.
- Educate the patient on how diabetes can affect the kidneys and provide tips on preventing damage to their kidneys:
- Controlling High Blood Pressure
- Medication Adherence by taking prescribed medication that protects the kidney functionality (ACE inhibitors or ARBs)
- Offer education on harmful medications to the kidneys (NSAIDS such a naproxen or ibuprofen)
- Suggest a diet of lower protein and limited salt intake
- Coordinate patient care with specialists (endocrinologist or nephrologist) as needed.

STATIN THERAPY FOR PATIENTS WITH DIABETES (SPD)

The percentage of members 40–75 years of age during the measurement year with diabetes who do not have clinical atherosclerotic cardiovascular disease (ASCVD) who met the following criteria:

Two rates are reported:

- 1. Received Statin Therapy: Members who were dispensed at least one statin medication of any intensity during the measurement year.
- 2. Statin Adherence 80%: Members who remained on a statin medication of any intensity for at least 80% of the treatment period

Improving HEDIS Measure

- Educate patients on the importance of statin medication adherence.
- Adherence to the SPD measure is determined by the member remaining on their
- prescribed high or low-intensity statin medication for 80% of their treatment period.
- Adherence is determined by pharmacy claims data (the plan will capture data each time the member fills their prescription).
- Use telehealth to manage medication adherence when applicable, this helps with consistency and continual medication adherence compliance.
- If the member's pharmacy benefit allows for a 90-day supply, write a prescription for 90- days.
- Refer to website for pharmacy formulary and coverage.

Links:

Peach State Health Plan: www.pshp.com

Ambetter: Provider Resources

Ambetter from Peach State Health Plan is underwritten by Ambetter of Peach State Inc., which is a Qualified Health Plan issuer in the Georgia Health Insurance Marketplace. This is a solicitation for insurance. © 2024 Ambetter of Peach State Inc. All rights reserved.

PSHP_060424_0246, AMB24-GA-HP-00064, GA-7884